

Phone: (07) 3280 3000  
3 Mary Street, BLACKSTONE, QLD 4304  
BARMAC INDUSTRIES PTY LTD



- Multi site of activity
- Water based
- Highly effective
- Excellent residual activity
- Broad spectrum
- Low cost

program  
Fungicide  
Preventative

# Fungol 500

New improved  
formulation

Consider

## Other protectors in the Barmac Range

- Fungol 500
- Rhysol DG
- Chloroturf DG
- Scurol DG
- Thiram
- Mancozeb DG

## The Barmac Protectors



BARMAC INDUSTRIES PTY LTD (Est 1955)


3 Mary Street, Blackstone, Queensland 4304 Phone: (07) 3280 3000

[www.barmac.com.au](http://www.barmac.com.au)

## DIRECTIONS FOR USE

SITUATION	DISEASE	STATE	RATE	CRITICAL COMMENTS
Ornamentals	Botrytis Blight ( <i>Botrytis cinerea</i> )	All States	100mL/100L water (Apply at minimum of 1.0L/ha)	Spray at 14 day intervals commencing when the disease first becomes apparent and continuing until conditions no longer favour the disease. Spraying Saintpaulia and Poinsettia flowers may result in some petal scorch. Tepid water should be used and wet plants protected from direct sunlight. This use is subject to an AFRAC fungicide anti-resistance strategy: Do not apply more than 3 consecutive Barmac Fungol 500 Fungicide or related dicarboximide sprays.
Recreational Turf	Brown Patch ( <i>Rhizoctonia solani</i> ) Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	Qld, NSW, Tas, SA, WA only	60 to 90mL/ 10 to 150L water/100m <sup>2</sup>	<b>Curative Program:</b> Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure. <b>For Preventative Treatment:</b> Apply as a monthly spray program. This use is subject to an AFRAC anti-resistance strategy. To reduce the possibility of resistance build-up to Barmac Fungol 500 Fungicide do not apply more than 2 consecutive sprays of Barmac Fungol 500 Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Brown Patch ( <i>Curvularia spp.</i> ) ( <i>Dreschlera spp.</i> ) ( <i>Fusarium culmorum</i> ) ( <i>Rhizoctonia solani</i> )	Vic only		
	Fusarium Patch ( <i>Fusarium nivale</i> )	NSW, Vic, Tas, SA, WA only	90mL/10 to 150L water /100m <sup>2</sup>	<b>Curative Program:</b> Repeat treatment after 14 days. <b>For Preventative Treatment:</b> Apply as a monthly spray program. This use is subject to an AFRAC anti-resistance strategy. To reduce the possibility of resistance build-up to Barmac Fungol 500 Fungicide do not apply more than two consecutive sprays of Barmac Fungol 500 Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Leaf Spot ( <i>Dreschlera spp.</i> ) Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	Qld, NSW, Vic, SA, WA only	60mL/10 to 150L water /100m <sup>2</sup>	
	Spring Dead Spot ( <i>Leptosphaeria spp.</i> )*	NSW, SA only	65mL per 10 to 150L water /100m <sup>2</sup>	<b>For Preventative Treatment:</b> Apply as a monthly soil drench throughout the year. Water in thoroughly after application. *When used in a preventative program this product will limit the development of Leptosphaeria and improve the rate of turf recovery. This use is subject to an AFRAC anti-resistance strategy. To reduce the possibility of resistance build-up to Barmac Fungol 500 Fungicide do not apply more than two consecutive sprays of Barmac Fungol 500 Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Spring Dead Spot ( <i>Helminthosporium spp.</i> , <i>Curvularia spp.</i> )	Vic, WA only		
	Helminthosporium disease affecting leaf, crown and roots: (Black and white Helminthosporium spot: <i>Bipolaris spp.</i> , <i>Drechslera spp.</i> , <i>Exserohilum spp.</i> )	Qld, NSW, Vic, SA, WA only	45mL/10 to 150L of water/100m <sup>2</sup>	<b>For Curative Treatment:</b> Apply as a foliar spray and repeat application after 7 to 14 days. For white spot symptoms on couch grass ( <i>Cynodon dactylon</i> ) only, use higher water volumes of 100 to 150L of water per 100m <sup>2</sup> . <b>For Preventative Treatment:</b> Apply as a high volume foliar spray. Repeat applications at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. To reduce the possibility of resistance build-up to Barmac Fungol 500 Fungicide do not apply more than two consecutive sprays of Barmac Fungol 500 Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION



For further information, or to place an order,  
please call

### LOW COST

Compare the cost of FUNGOL 500 with other fungicide applications, and you will see it makes it a viable alternative as a preventative program. In some instances it is less than half the cost of some of the other leading brands.

### BROAD SPECTRUM

FUNGOL 500 is extremely effective against a wide spectrum of damaging turf fungus which is an advantage in a preventative strategy.

### EXCELLENT RESIDUAL ACTIVITY

The improved formulation of FUNGOL 500 provides long term protection. Use in a regular program of 14 to 30 day applications.

### HIGHLY EFFECTIVE

FUNGOL 500 is extremely effective against some of the most difficult to control turf diseases including Brown Patch, Fusarium, Spring Dead Patch, Black and White Helminthosporium and Dollar Spot.

### MULTI SITE ACTIVITY

An advantage of Fungol 500 is its multi sited fungicidal activity. It interferes with the biological processes such as mycelium growth and cell division.

### WATER BASED

Fungol 500 is a water based fungicide with the flexibility of both curative and preventative actions. It is a low hazard to users, the environment and to the public.

Available in convenient to use 5 Litre containers.