



# POISON

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

▲ PEEL HERE ▲



# Pronamide

## Selective Herbicide

**ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE  
SOLVENT: 42 g/L ETHYLENE GLYCOL**

**GROUP D HERBICIDE**



**For use on Common Couch, Queensland Blue Couch, Kikuyu and Buffalo Grass.**

**For selective control of certain grasses and broadleaf weeds in lettuce and legume seed crops and pastures, as per the Directions for Use table.**

**IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING**



**Distributed by:** Amgrow Pty Ltd ABN 81 100 684 786

Unit B2A, 3-29 Birnie Avenue, Lidcombe NSW 2141 • TELEPHONE: 1800 631 008

CONTENTS

**5L, 10L, 20L**



# Pronamide

Selective Herbicide

## STORAGE & DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

## SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and waist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which can be obtained from the supplier.

**CONDITIONS OF SALE:** Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions.

APVMA Approval No.: 66067/58471

**IN A TRANSPORT EMERGENCY DIAL 000  
• POLICE OR FIRE BRIGADE •**



Batch Number

Date of Manufacture

[www.amgrow.com.au](http://www.amgrow.com.au)

## DIRECTIONS FOR USE

**Restrains:** DO NOT re-seed for 60 days after treatment.

DO NOT use on Bluegrass, Ryegrass, Fescue or Bent grass lawns.

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA	RATE BOWLING GREENS 37m x 37m	CRITICAL COMMENTS
Common Couch, Kikuyu	Winter Grass ( <i>Poa annua</i> )	All States	1.2 (12 mL/100m <sup>2</sup> )	170 mL	Apply during the autumn and winter, if established turf is infested with seeding Winter Grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25 mm irrigation or rain and again 24 hours later. Repeat if necessary on Qld Blue Couch after 4 weeks. Use only on established turf.
Queenstand Blue Couch		Qld only	0.6 (6 mL/100m <sup>2</sup> )	85 mL	
Buffalo Grass		WA only	1.2 (12 mL/100m <sup>2</sup> )	170 mL	

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA	CRITICAL COMMENTS
<b>Lettuce</b> Direct Sown	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Ryegrass,	All States	4.5 in 330-550 L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
Transplanted	Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	Vic only	2.3 in 300-550 L	Apply on light sandy soils, low in organic matter.

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA	CRITICAL COMMENTS
<b>Lucerne, Clover, Medics etc</b> Grown for forage, hay or seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats' Tail Fesuce	NSW, ACT, Tas, SA, WA only	1-1.5 L	<b>SEEDLING CROPS:</b> Apply after the third trifoliolate leaf is fully developed. <b>ESTABLISHED CROPS:</b> Apply any time before the end of July. <b>WEEDS:</b> Apply when weeds are post- or pre-emergent stages. <b>BEST RESULTS: When applied during cool and moist conditions.</b>
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, seedling Sorrel, Blackberry Nightshade, Paterson's Curse, seedling Wireweed		1.5-2 L	Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm of rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting <b>PRIOR</b> to treatment is recommended.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.**

#### **GENERAL INSTRUCTIONS**

Best results are obtained by applying the product to cool, wet soil.

#### **MIXING**

Prior to pouring, shake container vigorously, then add the required quantity of Pronamide Selective Herbicide to water in the spray vat while stirring or with agitators in motion.

DO NOT mix with other pesticides.

## RESISTANT WEEDS WARNING

Pronamide Selective Herbicide is a member of the benzamide group of herbicides. The product has the inhibitors of microtubule assembly mode of action. For weed resistance management Pronamide Selective Herbicide is a Group D herbicide. Some naturally-occurring weed biotypes resistant to Pronamide Selective Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Pronamide Selective Herbicide or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Amgrow Pty Ltd accepts no liability for any losses that may result from failure of Pronamide Selective Herbicide to control resistant weeds.

Advice as to strategies and alternative treatments that can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or Amgrow representative.

## PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT graze sheep on treated areas or feed to poultry or pigs.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, which may cause spray drift onto nearby susceptible plants, crops, cropping land or pastures.

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost. Triple rinse containers before disposal. Add rinsings to spray

GROUP **D** HERBICIDE

tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

## SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and waist, washable hat and elbowlength PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which can be obtained from the supplier.

**CONDITIONS OF SALE:** Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions.

\* Other trademarks

APVMA Approval Number: 66067/58471