

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



OxStar Herbicide Coated Fertiliser

ACTIVE CONSTITUENT: 10 g/kg OXADIAZON
SOLVENT: 10.5 g/kg POLYBUTENE

NPK 12 - 5 - 14

GUARANTEED FERTILISER ANALYSIS				w/w
Nitrogen (N) Total 12.0%	Phosphorus (P) Total 5.2%	Potassium (K) 14.1%	Sulfur (S) as Sulfate 8.0%	
Ammonium Form 7.0%	Water Soluble 2.6%	as Sulfate	Magnesium (Mg) as Sulfate 1.2%	
Nitrate Form 5.0%	Citrate Soluble 2.6%			
OxStar Herbicide Coated Fertiliser 12 - 5 - 14 is a versatile compound NPK fertiliser that provides uniform delivery of targeted, balanced nutrition to promote vigour in turf.				

Combination Herbicide and Controlled Release Fertiliser for Pre-Emergent Control of Summer Grass, Crab Grass and Winter Grass in Turf. Plus Nutrient Delivery (Feeding) to Turf.

Please Note: OxStar Herbicide is a pre-emergent herbicide. It will not kill the weeds that are already present. It only stops weeds if applied before weed germination.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA Approval No.: 83464/135291HG



Turf Culture
 43 Gap Road
 Sunbury Vic 3429
 Ph: 1300 11 8873
 (Ph: 1300 11 TURF)
www.turfculture.com.au

HOW TO USE

Situation / Turf Species	Weeds Controlled	Rate	Critical Comments
Established Turf & Lawns species as listed: Bahia Grass (<i>Paspalum notatum</i>), Buffalo grass (<i>Stenotaphrum secundatum</i>), Common Couch (<i>Cynodon dactylon</i>), Hybrid Couch (<i>Cynodon dactylon</i> C. <i>transvaalensis</i>) [NOT Santa ana], Kikuyu (<i>Pennisetum clandestinum</i>), Queensland Blue Couch (<i>Digitaria didactyla</i>), Saltwater Couch (<i>Paspalum vaginatum</i>), Zoysia (<i>Zoysia matrella</i>)	Pre-emergent control of: Summer Grass (<i>Digitaria ciliaris</i>), Crowsfoot Grass (<i>Eleusine indica</i>)	3 kg per 100m ²	The product is ready to apply. No mixing or dilution is required. Apply product evenly to dry foliage and bare areas using a granular spreader able to give even coverage. Application must be made before weeds have begun to germinate. Established weeds will not be controlled. After application water in with at least 10 mm of sprinkler irrigation. Control of germinating weed seedlings can be expected for up to 10 weeks after application depending upon temperature, rainfall, soil type and other factors. If longer control is required a repeat application should be made 7 to 10 weeks after initial application. DO NOT mow treated area until product has been watered in and turf foliage has dried. Avoid application when daytime temperatures are consistently above 32°C. This product is not recommended for use on putting greens, low cut tees or bowling greens. DO NOT apply on turf containing Carpet Grass (<i>Axonopus</i> spp.), Centipede Grass (<i>Eremochloa ophiuroides</i>) and Dichondra unless some damage is acceptable.
	Pre-emergent Control of: Winter Grass (<i>Poa annua</i>), Creeping Oxalis (<i>Oxalis comiculala</i>)	4 kg per 100m ²	
	Turf Nutrition: Both rates of this product will provide plant nutrients	3 to 4 kg per 100m ²	

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

OxStar Herbicide Coated Fertiliser (hereinafter OXSTAR) is a convenient ready to use formulation for the control of certain annual grass weeds in turf and also for fertilisation of turf where nutrients are required. The product should only be used when both functions are required.

OXSTAR is safe to use on the **established** turf species listed in the Situation column of the HOW TO USE table.

OXSTAR is a selective pre-emergent surface applied herbicide that provides residual control of grass and broadleaf weeds in established turf. OXSTAR controls susceptible weeds by preventing growth and development of newly germinated weeds.

HOW TO PREPARE**Soil Preparation**

Areas to be treated should be free of established weeds. Remove trash (weed residue, prunings, grass clippings etc.) from the surface before applying the product.

Measure

Before application, measure the lawn area to be sprayed, and determine the amount of OXSTAR required.

Application

For a granular spreader, which can be calibrated for even output and capable of giving even application. Apply under calm conditions, thereby reducing drift to non-target areas and optimizing particle distribution for good weed control. Ensure turf area is free of leaves and other debris, mow the area if necessary so that the granules reach the soil surface. DO NOT over apply, especially under the branch spread of trees and shrubs. Avoid contact with flowers and shrubs.

After application water in with at least 10 mm of sprinkler irrigation. DO NOT rely on forecast heavy rain or storms for OXSTAR incorporation, as heavy rain can cause runoff. Runoff (surface water movement) caused by heavy rainfall or heavy irrigation is when the soil is receiving more water than it can absorb. Runoff or pooling will move any OXSTAR just applied away from where the OXSTAR was placed by the spreader resulting in poor weed control from where OXSTAR was moved (by runoff).

Replanting / Overseeding Interval

Ryegrass: Allow 4 months after last application to lapse before re-seeding.

Bent grass: Allow 5 months after last application to lapse before re-seeding.

Timings

Apply prior to weed emergence in early spring (summer weeds) or late summer to early autumn (Winter Grass, *Poa annua*).

OXSTAR has excellent efficacy on annual germinating weeds such as Winter Grass, Summer Grass, Crowsfoot Grass (as listed in the How to Use table). The timings for application is therefore before germination of these weeds.

PRECAUTIONS

DO NOT apply to turf which is not well-established.

DO NOT apply if heavy rain has been forecast within 48 hours.

DO NOT apply to waterlogged soil.

DO NOT apply granules when the turf foliage is wet.

DO NOT use on putting greens, low cut tees or bowling greens.

DO NOT apply to newly seeded turf areas.

DO NOT mow treated area until product has been watered in and turf foliage has dried.

Avoid application when daytime temperatures are consistently above 32°C.

DO NOT apply on turf containing Carpet Grass (*Axonopus spp.*), Centipede Grass (*Eremochloa ophiuroides*) and Dichondra unless some damage is acceptable.

DO NOT graze treated turf / lawn or feed turf / lawn clippings from any treated area to poultry or livestock.

DO NOT use grass clippings from treated area for compost, around edible and non-edible plants or trees.

DO NOT allow containers or product to get into drains, sewers, ponds or streams.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated, dry place out of reach of children and pets. DO NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

Spills: Sweep up spilled materials and return to original container. Wash contaminated equipment taking care the washings DO NOT enter sewers, drains and other waterways.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. DO NOT inhale dust. Wash hands after use.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which is available from the supplier or from the manufacturer's website: www.turfculture.com.au

CONDITIONS OF SALE

The use of OXSTAR being beyond the control of the manufacturer, no warranty expressed or implied is given by Turf Culture regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Turf Culture accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

In a Transport Emergency Dial 000 Police or Fire Brigade